References

American Society of Anesthesiologists. (2024, January 29). The environmental impact of inhaled anesthetics. https://www.asahq.org/about-asa/governance-and-committees/asa-committees/environmental-sustainability/greening-the-operating-room/inhaled-anesthetics#:~:text=The%20atmospheric%20lifetimes%20of%20the,and%20sevoflurane%20(2%20years).

American Society of Anesthesiologists. (2023, October 18). Statement on the use of low gas flows for sevoflurane. https://www.asahq.org/standards-and-practice-parameters/statement-on-the-use-of-low-gas-flows-for-sevoflurane

Arslan, M., Gişi, G., Öksüz, G., Öksüz, H., Bilal, B., Boran, Ö. F., & Çalışır, F. (2020). Are high fresh gas flow rates necessary during the wash-in period in low-flow anesthesia? *The Kaohsiung Journal of Medical Sciences*, *36*(10), 834–840. https://doi.org/10.1002/kjm2.12251

Kitsiripant, C., Boonmuang, P., Chatmongkolchart, S., Tanasansuttiporn, J., Liochaichan, N., & Jantawong, N. (2022). A comparison of the cost and consumption of sevoflurane and litholyme as total gas flow rates of 1 and 2 liters per minute: a randomized controlled trial. *Journal of Health Science and Medical Research, 41*(1). doi: 10.31584/jhsmr.2022895 https://jhsmr.org/index.php/jhsmr/article/view/895/951

Kitson, A., Harvey, G., & McCormack, B. (1998). Enabling the implementation of evidence-based practice: a conceptual framework. Quality in Health Care, 7, 149-158. DOI: 10.1136/qshc.7.3.149. http://qualitysafety.bmj.com/content/7/3/149.abstract

Kutlusoy, S., Koca, E., & Aydin, A. (2022). Reliability of low-flow anesthesia procedures in patients undergoing laparoscopic cholecystectomy: Their effects on our costs and

ecological balance. *Nigerian Journal of Clinical Practice*, *25*(11), 1911–1917. https://doi.org/10.4103/njcp.njcp_387_22

McCormack, B., McCarthy, G., Wright, J., Slater, P., & Coffey, A. (2009). Development and testing of the context assessment index (CAI). *Worldviews on Evidence Based Nursing*, First Quarter, 27-35.

Shelgaonkar, V. & Choudhary, A. (2017). Low flow anesthesia: economic, eco-friendly, and effective. *Indian Journal of Clinical Anesthesia, 4*(3), 382-387. doi: 10.18231/2394-4994.2017.0077 https://www.ijca.in/article-download/full-text/4868

Simsek, T., Derman, S., Kordi, R., Saracoglu, A., & Saracoglu, K. (2022). The effect of different flow levels and concentration of sevoflurane during the wash-in phase on volatile agent consumption: a randomized controlled trial. *Journal of Clinical Monitoring and Computing, 36*, 1257-1262. https://doi.org/10.1007/s10877-022-00846-w

Singh, A., Sinha, R., Aravindan, A., Kumar, K., & Datta, P. (2019). Comparison of low-fresh gas flow technique to standard technique of sevoflurane induction in children: a randomized controlled trial. *Pediatric Anesthesia, 29*(4), 304-309. https://doi-org.proxy-hs.researchport.umd.edu/10.1111/pan.13582